

DIAS

All India TEST – SERIES CSE – 2024

Physical – 1

Syllabus :- Atomic structure, chemical bonding, liquid state, solid state, gaseous state

Instructions:-

1. Attempt five questions selecting atleast one question each section. apart from question 1&5 which is compulsory.
2. Write answer in space provided for this purpose only.
3. Total time allowed is 3hr and Marks is 250.

Information:

Name of student:- AAYUSH SAINI

UPSC Roll no:- 0841601

Mobile Number. [REDACTED]

Official use.

Q.NO.	1	2	3	4	5	6	7	8
MARKS	21	36	34	32	33			

Signature of invigilator

V. [Signature]

156
250
Signature of
Examiner

DIAS

All India TEST - SERIES CSE - 2024

Physical - I

Syllabus - Atomic structure, chemical bonding, liquid state, solid state, gaseous state

Instructions:-

1. Attempt five questions selecting atleast one question each section apart from question 1&2 which is compulsory.
2. Write answer in space provided for this purpose only.
3. Total time allowed is 3hr and Marks is 250.

Information:-

Name of student:- AAVUSH DAWI

UPSC Roll no:- 0841601

Mobile Number: [REDACTED]

Official use

Q.No.	1	2	3	4	5	6	7
MARKS	21	4	30	21	21	21	21

Signature of invigilator



Signature of
Examiner
122
080

DIAS

SECTION- A

1.(a) When an electron was accelerated through a potential difference of 1.00 ± 0.01 kilovolt, what is the uncertainty of position of the electron along its line of projection? (10)



(b). How far will the mercury level be depressed when a glass capillary with 0.40 mm. radius is placed in a dish of mercury? Given the surface tension of liquid mercury is 0.90 N/m and its density is $13.6 \times 10^3 \text{ kg m}^{-3}$.
(10)

When a capillary is dipped in Mercury, two forces F_{cohesion} , F_{adhesion} acts.
Since in mercury $F_{\text{cohesion}} > F_{\text{adhesion}}$,
Mercury level falls in capillary



Considering, h = height mercury falls
 r = radius of capillary = 0.4 mm
 γ = surface tension = 0.9 N/m
 ρ = density = $13.6 \times 10^3 \text{ kg/m}^3$
 θ = angle of contact
 Equating - Upward force due to surface tension = weight of the column goes down

$$\gamma(2\pi r \cos \theta) = \pi r^2 h \rho g$$

$$h = \frac{2\gamma \cos \theta}{\rho g r}$$

Considering $\theta = 180^\circ$ (as mercury level falls)

$$h = -\frac{2\gamma}{\rho g r}$$

$$h = \frac{-2(0.90 \text{ N/m})}{(13.6 \times 10^3 \text{ kg/m}^3)(9.8 \text{ m/s}^2)(0.4 \times 10^{-3} \text{ m})}$$

$$= -3.37 \times 10^{-2} \text{ m} = -3.37 \text{ cm}$$

Thus the mercury level drops by

$$3.37 \text{ cm}$$

DIAS

(c) The density of Li metal is 0.53 g cm^{-3} and the separation of the (100) planes of the metal is 350 pm . Determine whether the lattice is f.c.c or b.c.c $M(\text{Li}) = 6.941 \text{ g mol}^{-1}$ (10)

Since, wavelength of x-ray lies in range $10^{-10} - 10^{-12} \text{ cm}$, which is equivalent to size of an atom, it is used to determine the position of atoms.

Given that $(hkl) = (100)$
 $d(\text{separation of planes}) = 350 \text{ pm}$

$$\text{Let } a = \frac{\text{Side of a unit cell}}{d_{hkl}} = \frac{1}{\left(\frac{h}{a}\right)^2 + \left(\frac{k}{b}\right)^2 + \left(\frac{l}{c}\right)^2}$$

when $h, k, l =$ Miller indices
 $a = b = c$
 (for cube)

$$\therefore d_{hkl} = a = 350 \text{ pm} \quad \text{--- (i)}$$

Given that $\rho = 0.53 \text{ g/cm}^3$ (density)
 $M = 6.941 \text{ g/mol}$ (molar mass)
 $Z = \text{number of atoms per unit cell}$



we know, $\rho = \frac{ZM}{N_0 a^3}$ where $N_0 = 6.022 \times 10^{23} / \text{mol}$

$$(0.53 \text{ g/cm}^3) = \frac{Z (6.941 \text{ g/mol})}{(6.022 \times 10^{23} / \text{mol}) (350 \times 10^{-10} \text{ cm})^3}$$

$$Z \approx 1.97$$

$$\boxed{Z \approx 2}$$

Thus, Li is arranged in **BCC** lattice




DIAS

(d) At what pressure does the mean free path of argon at 25°C become comparable to the diameter of the atoms themselves? Given $\sigma = 0.36\text{ nm}^2$. (10)

Mean free path is the total distance travelled by the molecule in motion between two subsequent collisions.

Let us take a molecule of diameter σ , travelling with a average velocity $\bar{c}$.



Thus the molecule 'A', to prevent collision draws a cylinder of radius σ .
 Volume swept/sec = $\pi \sigma^2 \bar{c}$

No. of molecules undergoing collision = $\pi \sigma^2 \bar{c} N_A$

Taking relative motion consideration [where $N_A = \frac{N}{\text{Volume}}$]
 Z_1 (collision number) = $\sqrt{2} \pi \sigma^2 \bar{c} N_A$

Since λ (mean free path) = $\frac{\bar{c}}{Z_1} = \frac{1}{\sqrt{2} \pi \sigma^2 N_A}$ — (i)

Assuming ideal gas behaviour

$$PV = \frac{N}{N_A} RT \Rightarrow P = N_A K T \quad \text{--- (ii)}$$

$$N_A = \frac{P}{K T} \quad \text{--- (iii)}$$

Using equation (iii) in (i)

$$\lambda = \frac{K T}{\sqrt{2} \pi \sigma^2 P}$$

$$P = \frac{K T}{\sqrt{2} \pi \sigma^2 \lambda}$$

Given $\sigma = 0.36\text{ nm}$

$$P = \frac{(1.38 \times 10^{-23}) (298\text{ K})}{\sqrt{2} \pi (0.36 \times 10^{-9})^2 (0.36 \times 10^{-9})}$$

$$= 1985.2 \times 10^4 \text{ Pascals}$$

$$P = 198.5 \text{ bar}$$

(e) Explain on the basis of MO theory, why:

- Liquid oxygen sticks to the poles of a magnet while liquid nitrogen does not do so.
- He_2 and Be_2 molecules do not exist.
- The bond order in oxygen is 2, while that of nitrogen is 3.
- Hydronium molecule is more stable than H_2^+ ion.

(10)

MO Theory is based on mixture of all the orbitals of atoms to form equal number of Bonding and Antibonding orbital. The theory is provided by Pauling.


(i) Magnetic property depends on number of unpaired electrons using MOT.

$$\text{Bond Order} = \frac{N_{\text{BMO}} - N_{\text{ABMO}}}{2}$$

$$= \frac{4 - 4}{2} = 0$$

For He_2

$$1s^2$$



$$\text{Bond order} = \frac{2 - 2}{2} = 0$$

$\therefore$ Both have Bond order = 0

(ii) From MOT diagram of part (i)


$$\text{B.O.}(\text{O}_2) = \frac{8 - 4}{2} = 2$$

$$\text{B.O.}(\text{N}_2) = \frac{8 - 2}{2} = 3$$

Thus O_2 is Paramagnetic and sticks to poles and N_2 is Diamagnetic, does not do so.


(iii) Be_2

$$(\text{Be} = 1s^2 2s^2)$$



(iv)

$$1s^1$$

 H_2 

$$\text{B.O.}(\text{H}_2) = \frac{2 - 0}{2} = 1$$

$$\text{For } \text{H}_2^+ \quad \text{B.O.} = \frac{1 - 0}{1} = 0.5$$

Thus $\text{B.O.}(\text{H}_2) > \text{B.O.}(\text{H}_2^+)$

H_2 is more stable

2.(a) An atom makes a transition from an excited state with lifetime of 1 ns to the ground state by emitting a photon of wave length 600 nm. Calculate the uncertainty in the energy of the excited state. Also calculate the percentage uncertainty if the energy is measured from the ground state. (20)

3(a) Derive virial equation of state starting from Vanderwall equation of state.

(10)

Vanderwaal equation is given by
(for real gases)

$$\left(P + \frac{a}{V_m^2}\right)(V_m - b) = RT \quad \text{---(i)}$$

where,

P = Pressure

a = Pressure correction factor

 V_m = Molar volume

b = Volume correction factor

R = Gas Constant

T = Temperature.

$$\left(P + \frac{a}{V_m^2}\right) = \left(\frac{RT}{V_m - b}\right)$$

$$P = \frac{RT}{V_m - b} - \frac{a}{V_m^2}$$

$$\frac{PV_m}{RT} = Z = \frac{V_m}{V_m - b} - \frac{a}{V_m RT}$$

where

Z = Compressibility factor

$$Z = \left(\frac{1}{1 - b/V_m}\right) - \frac{a}{V_m RT}$$

Since $b \ll V_m$

$$\therefore Z = \left(1 - b/V_m\right)^{-1} - \frac{a}{V_m RT}$$

$$= \left[1 + \frac{b}{V_m} + \left(\frac{b}{V_m}\right)^2 + \left(\frac{b}{V_m}\right)^3 + \dots\right] - \frac{a}{V_m RT}$$

$$= 1 + \frac{b}{V_m} + \left(b - \frac{a}{RT}\right) \frac{1}{V_m^2} + \frac{b^2}{V_m^3} + \dots$$

$$Z = 1 + \frac{B}{V_m} + \frac{C}{V_m^2} + \frac{D}{V_m^3} + \dots$$

where

B, C, D. - virial coefficients

where $B = b - \frac{a}{RT}$, $C = b^2$,
 $D = b^3$ and so on

Further, virial equation can

be utilized to derive Boyle's Temp.

(Temp. at which Real gas behaves as ideal)

$$\text{With } B = 0 = b - \frac{a}{RT} \Rightarrow$$

$$T = \frac{a}{Rb}$$

Considering higher order
Coeff. negligible.


(b) Give the schrodinger's wave equation for H-atoms in cartesian and polar co-ordinates. With the help of a diagram show the relation between the two coordinates. (10)

In Cartesian Coordinates

$$\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right) \psi + \frac{8\pi^2 m}{h^2} (E - V) \psi = 0$$

When

ψ = wave function
 m = mass of molecule
 E = Total Energy
 V = Potential Energy



Correlation:

$$z = r \cos \theta$$

$$x = r \sin \theta \cos \phi$$

$$y = r \sin \theta \sin \phi$$

{ provide range }
 $0 < r < \infty$
 $0 < \theta < \pi$
 $0 < \phi < 2\pi$

$$\tan \phi = y/x$$

Schrodinger Equation in Polar coordinates

$$\psi = R \Theta \Phi$$

$$\frac{1}{R} \frac{\partial}{\partial r} \left(r \frac{\partial R}{\partial r} \right) + \frac{1}{\Theta \sin \theta} \frac{\partial}{\partial \theta} (\Theta \sin \theta) + \frac{1}{\Phi \sin^2 \theta} \frac{\partial^2 \Phi}{\partial \phi^2} + \frac{8\pi^2 m}{h^2} (E - V) = 0$$

From this ψ can be derived which is used to determine the position of wave function.

DIAS

(c) In a certain material of simple cubic structure, (100) diffraction is obtained at $\theta = 24.88^\circ$ with radiation of $\lambda = 1.541 \text{ \AA}$. Can this material accommodate an atom of $1.68 \text{ \AA}$ radius interstitially in void space without lattice distortion? (10)

Since, X-ray wavelength lies in range $10^{-10} - 10^{-12} \text{ m}$, it is utilized to determine lattice structure. Formula for which is provided by

Bragg's equation

$$n\lambda = 2d \sin \theta$$

where λ = wavelength
 d = distance between planes

θ = Angle of reflection
 n = order of reflection


determining d_{hkl} ,

$$\lambda = 2 \left(\frac{d}{n} \right) \sin \theta$$

$$\lambda = 2 d_{hkl} \sin \theta$$

$$d_{hkl} = \frac{\lambda}{2 \sin \theta} = \frac{(1.541) \text{ \AA}}{2 (\sin 24.88^\circ)}$$

$$d_{hkl} = 1.83 \text{ \AA}$$



Since

$$d_{hkl} = \frac{a}{\sqrt{h^2 + k^2 + l^2}}$$

a = side of a unit cell
 h, k, l = miller indices

$$d_{hkl} = \frac{a}{\sqrt{1+0+0}}$$

$$\Rightarrow a = 1.83 \text{ \AA}$$

Finding size of interstitial void

Since

$$\frac{\sqrt{3}a}{2} = 2r_a = a$$

$$r_a = \frac{a}{2} = 0.915 \text{ \AA}$$

$$\text{Also, } \frac{\sqrt{3}a}{2} = r_a + r_v \Rightarrow r_v = \frac{\sqrt{3}}{2} (1.83) - (0.915)$$

$$r_v = 0.67 \text{ \AA} = \text{radius of void}$$

$$\text{Since } r_v < 1.68 \text{ \AA}$$

(It cannot be accommodated)

Provide for redrawing notes also

DIAS INDIA EDUTECH (P)LTD.

18, Pusa Road Karol Bagh, New Delhi -110005

PH: 011-40079000, 9350934622

(d) Using VSEPR theory, predict the shapes of PCl_2F_3 and SF_4 . Indicate the state of hybridization in each case. (10)

VSEPR theory helps in determining the shapes and geometry of the hybrid orbitals discussed by Valence bond theory (VBT)

It uses, Drago's Formula = $\frac{1}{2} (V + M - A + C)$
(Bond Pair) (B.P.) where


V = Valency of Central atom
 M = Number of monatomic atoms
 A = Anionic charge
 C = Cationic charge

(i) PCl_2F_3

$$\text{B.P.} = \frac{1}{2} (5 + 5) = 5$$

$\therefore$ Hybridization = $\boxed{sp^3d^2}$

Structure



Trigonal Bi-Pyramidal
($\mu \neq 0$)

Since, Bent Rule states that electronegative atom goes to position of less % s-character. 2-F atoms become axial


(ii) SF_4

$$\text{B.P.} = \frac{1}{2} (6 + 4) = 5$$

$\boxed{sp^3d^2}$

$\Rightarrow$ Hybridization

$$\text{Lone Pair} = \text{B.P.} - 4 = 1$$



Shape = Saw
Geometry = Trigonal Bi-Pyramidal

Since, lone pair occupies more volume, it remains at equatorial position, where angle is 120°

(e) The energy levels of the electron in hydrogen atoms are given by

$$E_n = -R_H / n^2, n = 1, 2, 3, \dots$$


Where R_H is the Rydberg constant and n is the principal quantum number.

- Draw a clearly labelled energy level diagram showing the first three energy levels of the electron and all of the quantum states belonging to each energy level.
- Give the degeneracies of the first three energy levels, giving the allowed l and m_l values and the orbitals rotations. Suggest a simple formula which relates the degeneracy of an energy level to its principal quantum number. (10)

Quantum numbers are provided by Schrodinger wave equation solutions

Such that

Principal quantum number (n) → Determines energy
 Azimuthal quantum number (l) → Angular momentum
 Magnetic quantum number (m) → Direction of Angular momentum along z-direction



(ii) Degeneracies are the energy equivalent states. Since, energy depends only on ' n ', and is independent of ' l ' and ' m ', different l and m positions have same energy belonging to particular shell (n).

Since, $l = n-1$ (maximum value)
 and $m = 2l+1$ (no. of values)

$$\therefore \text{Degeneracy} = \sum_{l=0}^{n-1} (2l+1)$$

$$(D) \quad D = 2 \sum_{l=0}^{n-1} l + \sum_{l=0}^{n-1} 1 \Rightarrow \frac{2 \times (n-1)(n)}{2} + n$$





$$D = n^2$$

$\therefore$ No. of degeneracy of an energy level is n^2

4(a) Set up Born-Haber cycle for determining the enthalpy of solvation of Mg^{2+} ions from MgCl_2 by water.

(10)


(d) Which of the following pairs is expected to have the larger bond angle and why? (10)



SECTION - B

5(a) Derive the packing fraction for HCP.

(10)
HCP is a specific form of FCC (face centered cubic) such that ÷



Packing fraction (v) = $\frac{\text{Volume of all atoms } (V_a)}{\text{Volume of the trapezium } (V_t)} \quad \text{--- (X)}$

$$V_a = 6 \times V_{\text{atom}} = 6 \times \frac{4}{3} \pi r^3 \quad \text{where } r = \text{radius of atom}$$

$V_t \Rightarrow$ Assuming, most compact form (111) of FCC is equivalent in HCP.

$$\therefore \left[\frac{c}{2} = \frac{a}{\sqrt{3}} \right] \Rightarrow c = \frac{2a}{\sqrt{3}} \quad \text{--- (i)} \quad \left\{ a = \text{interplanar distance} \right\}$$

Also, for FCC, $4r = \sqrt{2}a$ --- (ii)

$$\therefore \text{Using (i) and (ii)} \quad \left[r = \frac{\sqrt{6}}{8} c \right]$$

$$\therefore V_t = 6 \times A_{eq} \times c \quad \left(\text{where } A_{eq} = \text{Area of equilateral triangle} \right)$$

$$\Rightarrow \sqrt{3} \pi r^2$$

$$V_t = 6 \times \sqrt{3} \pi r^2 \times \frac{8r}{\sqrt{6}} \quad \text{--- (iv)}$$

Putting V_a and V_t in eq (X).

$$\eta = \frac{6 \times \frac{4}{3} \pi r^3}{6 \times \frac{\sqrt{3}}{2} \times \frac{8r}{\sqrt{6}}} = \frac{\pi}{\sqrt{18}} = 0.7401$$

$$\therefore \boxed{\eta = 74.01\%}$$

Eg. of such packing are Hg and Cd

(b) What is Stoke's law. How it is utilized to determine viscosity of liquid?

(10)

Stoke's law is based on equalization of forces, such that terminal velocity is achieved by a solid steel ball falling through a liquid column. It states drag force to be

$$F_d = 6\pi\eta r v \quad \text{---(i)}$$

for terminal velocity,

$$F_d = F_g$$

F_d = force of Drag.

F_g = gravitational force

$$F_d = \frac{4}{3}\pi r^2 (\rho_s - \rho_l) g$$

$$F_g = \frac{4}{3}\pi r^3 \rho_s g$$

$$F_g = \frac{4}{3}\pi r^3 (\rho_s - \rho_l) g \quad \text{---(ii)}$$

where η = coeff. of viscosity
 r = radius of ball
 v = Terminal velocity
 ρ_l = density of liquid
 ρ_s = density of ball
 $g = 9.81 \text{ m/s}^2$.

Equating (i) = (ii)

$$\frac{3}{2} 6\pi\eta r v = \frac{4}{3}\pi r^3 (\rho_s - \rho_l) g$$

$$\eta = \frac{2}{9} \frac{r^2 (\rho_s - \rho_l) g}{v}$$

$$v = \frac{l}{t} \Rightarrow \eta = \frac{2}{9} \frac{r^2 (\rho_s - \rho_l) g t}{l}$$

where l = length
 t = time

It can be utilized as,

$$\frac{\eta_l}{\eta_w} = \frac{(\rho_s - \rho_l) t_l}{(\rho_s - \rho_w) t_w}$$

to determine unknown viscosity using known one.

(c) Verify that the wave function $\psi(x) = x e^{-ax^2}$ is an eigenfunction of the operator $d^2/dx^2 - 4a^2x^2$. What is the corresponding eigenvalue? (10)

Wave function when operates over an operator and the function gets regenerated along with a constant value, is called as an eigenfunction of that operator.

$$\hat{A} \psi = E \psi$$

Operator $\downarrow$ Eigen function $\rightarrow$ Eigen value

$$\hat{A} = \frac{d^2}{dx^2} - 4a^2x^2$$

$$\hat{A} \psi = \left(\frac{d^2}{dx^2} - 4a^2x^2 \right) (x e^{-ax^2})$$

$$= \frac{d^2}{dx^2} (x e^{-ax^2}) - 4a^2x^3 e^{-ax^2}$$

$$= \frac{d}{dx} (x e^{-2ax} - 2ax e^{-2ax}) - 4a^2x^3 e^{-2ax}$$

$$= -2a x e^{-2ax} - 2a e^{-2ax}$$

$$\hat{A} \psi = \frac{d}{dx} (x e^{-ax^2} - (ax)(2x) e^{-ax^2}) - 4a^2x^3 e^{-ax^2}$$

$$\hat{A} \psi = (-a)(2x) e^{-ax^2} - (2ax^2)(-2ax) e^{-ax^2} - 4a^2x^3 e^{-ax^2}$$

$$= -6ax e^{-ax^2}$$

$$\hat{A} \psi = -6 \psi$$

Thus ψ is a eigen function of operator $\frac{d^2}{dx^2} - 4a^2x^2$ with Eigen

$$\text{value } \boxed{-6}$$

(d) ((i) Construct a trial LCAO – MO wavefunction for the H_2 molecule. (10)

ii) The trial VB wavefunction used by Heitler and London to describe the H_2 molecule in terms of hydrogen – like wavefunction is

$$\phi = c_1 \psi_a(1) \psi_b(2) + c_2 \psi_a(2) \psi_b(1)$$

Compare with the wavefunction you written in (i) and comment on the differences, if any.

(i) For H_2

$$\psi_A = c_A \psi_A$$

$$\psi_B = c_B \psi_B$$


$$\psi_A - \psi_B \Rightarrow \boxed{\psi = c_A \psi_A \pm c_B \psi_B}$$

Bonding $\Rightarrow \psi = c_A \psi_A + c_B \psi_B$
M.O.

Antibonding $\Rightarrow \psi = c_A \psi_A - c_B \psi_B$
M.O.

(ii) $\phi = c_1 \psi_a(1) \psi_b(2) + c_2 \psi_a(2) \psi_b(1)$

~~This considers resonance~~




~~This considers exchange of e^- between the two hydrogens while above one doesn't.~~

(e) Write down equation for distribution of molecular speed on the basis of this explains the temperature dependence of distribution. (10)

Since, gaseous molecules are under continuous elastic collision, continuous kinetic exchange takes place, ~~it thus~~, rather than speed of individual atom, distribution is studied

Maxwell-Boltzmann Equation of molecular speed distribution

$$\frac{1}{N} \frac{dN}{dc} = 4\pi c^2 \left(\frac{m}{2\pi KT} \right)^{3/2} \exp\left(-\frac{mc^2}{2KT}\right)$$




Where, m = mass of molecule
 $K = 1.38 \times 10^{-23}$ (Boltzmann constant)
 c = speed of molecule
 $\frac{dN}{N} =$ fraction of molecule at particular speed.
 T = Temperature (in K)

Dependence of Temperature $\Rightarrow$

$$c \propto \sqrt{\frac{1}{T}}$$

$$T_1 < T_2 < T_3$$

As temperature increases, fraction of molecules associated with higher speed increases, while fraction of molecules with lower speed decreases. (Since total no. of molecules remain const.)



& dist. ^{become more} broader.

7(a) the density of ethyl acetate at 30°C is 0.9850 g cm^{-3} . Calculate the surface tension of ethyl acetate, given the parachor values of C, H, O and the double bond are 8.2, 16.3, 19.5 and 21.5 respectively. (10)

Eotvos gave Temperature dependence of surface tension
 $\gamma \frac{M_m}{\rho} r^{2/3} = a + bt$

Further Ramsay's modification $\frac{M_m}{\rho} r^{2/3} = b(t - t_c + b)$
 Further McLeod gave the concept of Parachor

Such that,

$$\boxed{\frac{M_m}{\rho} \gamma^{1/4} = P} \quad \text{--- (i)}$$


where

M_m = molar mass

ρ = density

γ = surface tension

P = Parachor values



Calculating P

$$4P_C + 2P_O + 8P_H + 1P_{DB}$$

$$= (4)(8.2) + 2(19.5) + 8(16.3) + 21.5$$

$$\boxed{P = 223.7}$$

$$M_m = 88 \text{ g/mol}$$

$$\rho = 0.9850 \text{ g/cm}^3$$

Putting in (i)

$$\frac{88}{0.9850} \gamma^{1/4} = 223.7$$

$$\boxed{\gamma = 39.307 \text{ dyn/cm}}$$

(Parachor is additive property)

(b) Show that Hermitian operator always gives real eigen value whenever applied to normalized wave function. (10)

Hermitian operator $\Rightarrow$

$$\langle \Psi_n^* | A | \Psi_n \rangle = \langle \Psi_n^* | A^* | \Psi_n \rangle$$

such that $A =$ ~~eigen~~ operator
 $\Psi_n =$ wave function

we know,

$$\cancel{\Psi_n^* A}$$

$$\hat{A} \Psi_n = a_n \Psi_n \text{ (i) when}$$

$$\hat{A}^* \Psi_n^* = a_n^* \Psi_n^* \text{ (ii) } \Psi_n^* = \text{Complex Conjugate of } \Psi_n$$

$a_n =$ Eigen values
 $a_n^* =$

Multiplying eq. (i) by Ψ_n^*
 and eq. (ii) by Ψ_n

$$\Psi_n^* \hat{A} \Psi_n = a_n \Psi_n^* \Psi_n \text{ (i)}$$

$$\Psi_n \hat{A}^* \Psi_n^* = a_n^* \Psi_n \Psi_n^* \text{ (ii)}$$

Subtracting and integrating over space.

$$\int_{\text{space}} (\Psi_n^* \hat{A} \Psi_n - \Psi_n \hat{A}^* \Psi_n^*) dZ = \int_{\text{space}} (a_n - a_n^*) \Psi_n \Psi_n^* dZ$$

$\rightarrow$ As $\hat{A}$ is hermitian operator

$$\boxed{LHS = 0}$$

$$\int_{\text{space}} (a_n - a_n^*) \Psi_n \Psi_n^* dZ = 0$$

$$\text{Since } \int_{\text{space}} \Psi_n \Psi_n^* dZ = 1$$


$$\boxed{a_n = a_n^*}$$

Thus, both real and complex eigen values are equal iff $(a_n = a_n^*) \Rightarrow$ both are real.

(c) Derive Heisenberg uncertainty principle for particle in 1-D box.

(10)

Assuming, a box of length l and particle moving along only the x -direction, $V=0$ (inside box)
 $V=\infty$ (outside)



As per Schrodinger wave eqⁿ

$$\psi = \sqrt{\frac{2}{l}} \sin\left(\frac{n\pi x}{l}\right)$$

Heisenberg uncertainty principle $\Delta x \cdot \Delta p_x \geq \frac{h}{4\pi}$

$$\Delta v = \sqrt{\langle v^2 \rangle - \langle v \rangle^2}$$

$$\therefore \langle \psi | \hat{p}_x | \psi \rangle = \int_0^l \left(\frac{2}{l}\right) \sin\left(\frac{n\pi x}{l}\right) (-i\hbar) \frac{\partial}{\partial x} \left(\sin\left(\frac{n\pi x}{l}\right)\right) dx$$

$$= \int_0^l \left(\frac{2}{l}\right) (-i\hbar) \left(\frac{n\pi}{l}\right) \sin\left(\frac{n\pi x}{l}\right) \cos\left(\frac{n\pi x}{l}\right) dx$$

$$\langle \hat{p}_x \rangle = 0$$

$$\langle \psi | \hat{p}_x^2 | \psi \rangle = \int_0^l \left(\frac{2}{l}\right) (-\hbar^2) \left(\frac{n\pi}{l}\right)^2 \sin^2\left(\frac{n\pi x}{l}\right) dx$$

$$\langle \hat{p}_x^2 \rangle = \frac{n^2 \hbar^2}{4l^2}$$

$$\langle \psi | \hat{u} | \psi \rangle = \int_0^l \left(\frac{2}{l}\right) x \sin^2\left(\frac{n\pi x}{l}\right) dx = \frac{l}{2}$$

$$\langle \hat{u} \rangle = \frac{l}{2}$$

$$\langle \psi | \hat{u}^2 | \psi \rangle = \int_0^l \left(\frac{2}{l}\right) x^2 \sin^2\left(\frac{n\pi x}{l}\right) dx = \frac{l^3}{12}$$

$$\langle \hat{u}^2 \rangle = \frac{l^3}{12}$$

$$\Delta \hat{p}_x = \sqrt{\langle \hat{p}_x^2 \rangle - \langle \hat{p}_x \rangle^2} = \frac{n\hbar}{2l}$$

$$\Delta \hat{u} = \sqrt{\langle \hat{u}^2 \rangle - \langle \hat{u} \rangle^2}$$

DIAS

(d) Discuss the structure of IF_7 and $POCl_3$ on the basis of hybridization.

V.B.T. along with USEPR theory explains the shape and structure of covalent molecules.

Such that, as per Drago's formula

$$\text{B.P. (Bond Pair)} = \frac{1}{2} (U + M - A + C)$$


U = Valency of Central atom

M = number of monovalent atoms

A = Anionic Charge

C = Cationic Charge

for,




$$\text{B.P.} = \frac{1}{2} (7 + 7) = 7$$

$$\text{L.P. (Lone pair)} = 0$$


Hybridization = sp^3d^3

Pentagonal bi-Pyramidal



(Regular shape)



sp^3



$$\text{B.P.} = \frac{1}{2} (5 + 3) = 4$$



Tetrahedral shape




(e) Draw the radial part of the wave function and radial probability distribution for hydrogen atoms for 1s, 2s and 2p orbitals. (10)

1s

$$R_{1s} = 2 \left(\frac{1}{a_0} \right)^{3/2} \exp \left(-\frac{r}{a_0} \right)$$

$$P_{1s} = 4\pi r^2 R_{1s}^2$$

$$4\pi r^2 \left[\frac{4}{a_0^3} \exp \left(-\frac{2r}{a_0} \right) \right]$$





when $a_0 = 0.53 \text{ \AA}$

2s

$$R_{2s} = \left(\frac{1}{2a_0} \right)^{3/2} \left(2 - \frac{r}{a_0} \right) \exp \left(-\frac{r}{2a_0} \right)$$



$$P_{2s} = 4\pi r^2 \left[\frac{1}{8a_0^3} \left(2 - \frac{r}{a_0} \right)^2 \exp \left(-\frac{r}{a_0} \right) \right]$$



2p

$$R_{2p} = \frac{1}{\sqrt{3}} \left(\frac{1}{2a_0} \right)^{3/2} \left(\frac{r}{a_0} \right) \exp \left(-\frac{r}{2a_0} \right)$$

$$P_{2p} = 4\pi r^2 \left[\frac{1}{3} \left(\frac{1}{8a_0^3} \right) \left(\frac{r^2}{a_0^2} \right) \exp \left(-\frac{r}{a_0} \right) \right]$$



DIAS

8(a). The critical temperature and pressure for CO_2 gas are 304 K and 74 atm, respectively, and for NH_3 are 405 K and 111 atm, respectively. Which gas has the smaller values of the van der Waals' constants, a and b ? Which is the most nearly ideal in behavior at 300 K and 10 atm? (10)

Critical temperature is the one above which no liquification of gas is possible, whatever may be the pressure applied.

For real gas,

$$\left(P + \frac{a}{V_m^2}\right)(V_m - b) = RT$$


solving eq. $(V_{m,c} - V_{m,i})^3 = 0$

It gives,

$$V_c = 3b$$

$$P_c = \frac{a}{27b^2}$$

$$T_c = \frac{8a}{27Rb}$$



For CO_2

$$T_c = 304\text{K}$$

$$P_c = 74\text{ atm}$$

$$304 = \frac{8a_{\text{CO}_2}}{27Rb_{\text{CO}_2}}$$

$$74 = \frac{a_{\text{CO}_2}}{27b_{\text{CO}_2}^2}$$

$$\frac{304}{74} = \frac{8b_{\text{CO}_2}}{R}$$

$$b_{\text{CO}_2} = \frac{74}{8} \frac{304R}{(74)(8)}$$

Similarly

$$a = \frac{T_c(27Rb)}{8} = \frac{(T_c)(27)(R)}{8}$$

For NH_3

$$405 = \frac{8a}{27Rb}$$

$$111 = \frac{a}{27b^2}$$

$$\frac{405}{111} = \frac{8b}{R}$$

$$b_{\text{NH}_3} = \frac{405R}{(8)(111)}$$

$$b_{\text{CO}_2} > b_{\text{NH}_3}$$

and $a_{\text{NH}_3} > a_{\text{CO}_2}$

(b) What is atomic radius of a cubic lattice? Find the atomic packing factor in, body-centred cubic system and face-centred cubic system. (10)

(i) In cubic lattice




$r = \text{radius}$

$$2r = a$$

So, atomic radius is half the side of a unit cell in simple cubic lattice.

In BCC $2r = \frac{\sqrt{3}}{2} a$

In FCC $2r = \frac{\sqrt{2}}{2} a$



Atomic Packing $\Rightarrow \eta = \frac{\text{Volume occupied by atoms}}{\text{Volume of lattice}}$

For BCC $\Rightarrow \frac{2 \times \frac{4}{3} \pi r^3}{\left(\frac{4}{\sqrt{3}}\right)^3 r^3}$

$= 68\%$

$\boxed{Z=2}$


For FCC $\Rightarrow \boxed{Z=4}$

$= \frac{2 \times \frac{4}{3} \pi r^3}{(2\sqrt{2})^3 r^3}$

$= 74.01\%$

(c) Derive Bragg's equation. Using X-ray powder pattern of a compound describe how do you calculate the spacing between the planes. (10)

Since $\lambda_{\text{x-ray}} = 10^{-10} - 10^{-12} \text{ m}$ which is similar to size of atom. It can be used to determine structure of lattice.



For constructive interference
 $[DEF - ABC] = n\lambda$

$$DEF - ABC = GE + EH = 2d \sin \theta$$


where
 n = order of reflection

Therefore,

$$n\lambda = 2d \sin \theta$$

Powder method to determine spacing

(Exposure made)



Using Bragg's equation

$$d = \frac{n\lambda}{2 \sin \theta}$$


can be

utilized to determine spacing.

(d) What do you understand by resonance energy? How it is calculated.

(10)

Resonance is a phenomena when any system where no single structure represents complete picture of the molecule, such that a hybrid structure is made and the other structures are called canonical structures.



Resonance Energy → It is the energy difference between the most stable canonical structure and hybrid structure.

It can be calculated using! (Determined experimentally)

→ Heat of hydrogenation method.

→ Bond dissociation energy method etc. (Thermodynamic method)

→ Spectroscopy.

(e) why viscosity of gas increases with increase in temperature while that of liquid decreases with temperature. (10)

Viscosity is the resistance offered by the slower layer of fluid to the faster one.

For liquid $\Rightarrow$ Assumed that the fluid particles reside in holes (hypothetical)

$$\eta = \eta_0 \exp\left(\frac{E}{RT}\right)$$

E = Activation energy
 T = Temp.

η_0 = Intrinsic viscosity

As T increases, molecules come out of the holes frequently, η decreases

For gases $\Rightarrow$ As an temperature increasing kinetic energy of the gas increases, frequent collision occur, viscosity increases

$$\eta = \frac{1}{3} m n \lambda \bar{c}$$

Since both λ and $\bar{c}$ (mean free path and average velocity) increases with increase in temperature.

η increases for gas.